


Urology Journal

Official Journal of the Urology and Nephrology Research Center and the Iranian Urological Association



Intracorporeal Tapering of the Ureter for Distal Ureteral Stricture Before Laparoscopic Ureteral Reimplantation



Tuberous Sclerosis Complex

4th Biannual Congress of Iranian Endourology and
Urolaparoscopy Society
February 24-26, 2011 Tehran, Iran

Urology Journal

Official Journal of the Urology and Nephrology Research Center and the Iranian Urological Association

DIRECTOR

Nasser Simforoosh, MD

*Professor and Chairman, Department of Urology
Shahid Beheshti University of Medical Sciences*

EDITOR-IN-CHIEF

Abbas Basiri, MD

*Professor, Department of Urology
Director, Urology and Nephrology Research Center*

ASSOCIATE EDITOR

Mohammad Reza Safarinejad, MD

Professor, Department of Urology, AJA University of Medical Sciences

SECTION EDITORS

Endourology and Stone Disease

Abbas Basiri, MD

Masoud Etemadian, MD

Laparoscopic Urology

Nasser Simforoosh, MD

Alireza Amin-Sharifi, MD

Pediatric Urology

Abdolmohammad Kajbafzadeh, MD

Pejman Shadpour, MD

Urological Oncology

Seyed Yousef Hosseini, MD

Kidney Transplantation

Gholamreza Pourmand, MD

Mohammad Ali Zargar Shoushtari, MD

Female Urology

Farzaneh Sharifiaghdas, MD

Sexual Dysfunction and Infertility

Mohammad Reza Safarinejad, MD

Reconstructive Surgery

Jalil Hosseini, MD

Darab Mehraban, MD

Cellular and Molecular Urology

Farzaneh Sharifiaghdas, MD

Clinical Pathology Case

Farzaneh Sharifiaghdas, MD

EXECUTIVE EDITOR

Elham Iran-Pour, MD

ASSISTANT EDITORS

Amir Ali Tavabi, MD

Ali Reza Khoshdel, MD, PhD

ADMINISTRATIVE STAFF

Mojgan Khoddam

Morteza Rahmatizadeh

EDITORIAL BOARD

SMK Aghamir, MD *IR Iran*

H Ahmadnia, MD *IR Iran*

M Al-Mousawi, MD *Kuwait*

A Basiri, MD *IR Iran*

F Daneshgari, MD *USA*

S Daneshmand, MD *USA*

MR Desai, MD *India*

R Ghavamian, MD *USA*

S Gholami, MD *USA*

MA Haberal, MD *Turkey*

AIA Hussain, MD *Kuwait*

LR Kavoussi, MD *USA*

RB Khauli, MD *Lebanon*

Z Kirkali, MD *Turkey*

R Mahdavi Zafarghandi, MD *IR Iran*

RS Malek, MD *USA*

M Marberger, MD *Austria*

AR Mehra, MD *IR Iran*

MR Nikoobakht, MD *IR Iran*

G Pourmand, MD *IR Iran*

S Rizvi, MD *Pakistan*

H Sadeghi-Nejad, MD *USA*

MR Safarinejad, MD *IR Iran*

AA Samadi, MD *USA*

P Shadpour, MD *IR Iran*

FAM Shaheen, MD *Saudi Arabia*

A Shamsa, MD *IR Iran*

N Simforoosh, MD *IR Iran*

SAM Ziaee, MD *IR Iran*

MA Zargar Shoushtari, MD *IR Iran*

ISSN. 1735-1308

eISSN. 1735-546X

Publisher. Urology and Nephrology Research Center (UNRC)

Editorial Office. UNRC, No 103, 9th Boustan St, Pasdaran Ave, Tehran, Iran. Tel: +98 21 2259 4204, Fax: +98 21 2259 4204

Page Setter. Mahdi Akbarzadeh, graphic_negareh@yahoo.com

Printer. Ettehad Press, Tabas St, Ferdosi St, Imam Khomeini Sq, Tehran, Iran. Tel: +98 21 6671 6733

BOARD OF CONSULTANTS

HR Abdi, MD *IR Iran*
M Abedinzadeh, MD *IR Iran*
A Ahanian, MD *IR Iran*
M Aliasgari, MD *IR Iran*
AR Aminsharifi, MD *IR Iran*
E Amini, MD *IR Iran*
H Arshadi, MD *IR Iran*
SA Asgari, MD *IR Iran*
N Aydilek, MD *Turkey*
S Bagci, MD *Germany*
MR Barghi, MD *IR Iran*
CK Chang, MD *Taiwan*
SH Cohanzad, MD *IR Iran*
F Dadkhah, MD *IR Iran*
B Delahunt, MD *Australia*
H Djaladat, MD *IR Iran*
T Dogu, MD *Germany*
A El-Assmy, MD *Egypt*
LA Fariña, MD *Spain*
AR Farshi, MD *IR Iran*
S Falahatkar, MD *IR Iran*
M Ghaderian, MD *IR Iran*
A Gözen, MD *Germany*
Wu Guojun, MD *China*
R Haghghatkhah, MD *IR Iran*

G Haidl, MD *Germany*
S Hajebrahimi, MD *IR Iran*
B Helpap, MD *Germany*
P Hes, MD *Mexico*
M Hiradfar, MD *IR Iran*
S Horenblas, MD *Amsterdam*
M Houshmand, MD *IR Iran*
C Kamil, MD *Turkey*
B Kazemi, MD *IR Iran*
M Khazaeli, MD *IR Iran*
A Kheradmand, MD *IR Iran*
AR Lashay, MD *IR Iran*
V Lokeshwar, MD *USA*
B López, MD *Spain*
M Mahdavi Mazdeh, MD *IR Iran*
R Mahdavi Zafarghandi, MD *IR Iran*
R Maghsoudi, MD *IR Iran*
S Mehrabi Sisakht, MD *IR Iran*
M Miner, MD *Australia*
S Momenzadeh, MD *IR Iran*
MK Moslemi, MD *IR Iran*
SH Mousavi-Bahar, MD *IR Iran*
GH Naderi, MD *IR Iran*
H Naseh, MD *IR Iran*
M Nikkar, MD *IR Iran*

MR Nikoobakht, MD *IR Iran*
A Nouralizadeh, MD *IR Iran*
A Ozhand, MD *IR Iran*
B Papaziogas, MD *Greece*
E Razeghi, MD *IR Iran*
H Rezvani, MD *IR Iran*
K Richards, MD *USA*
M Roghani, MD *IR Iran*
P Shadpour, MD *IR Iran*
S Shakeri, MD *IR Iran*
F Sharifiaghdas, MD *IR Iran*
N Simforoosh, MD *IR Iran*
M Soleimani, MD *IR Iran*
A Tadayon, MD *IR Iran*
S Teber, MD *Turkey*
M Teshfam, MD *IR Iran*
P Tousi, MD *IR Iran*
MR Vaez-Mahdavi, MD *IR Iran*
C Valentina, MD *Italy*
M Yang B, MD *USA*
S Zare, MD *IR Iran*
M Zargham, MD *IR Iran*
A Zolfaghari, MD *IR Iran*

INTERNATIONAL ADVISORY BOARD

SA Adebaya, MD *Nigeria*
M Al-Mousawi, MD *Kuwait*
H Al-Zahrani, MD *Saudi Arabia*
AB Babanoori, MD *USA*
M Beheshti, MD *Canada*

R Davis, MD *USA*
MA Elakkad, MD *Egypt*
M Eshghi, MD *USA*
R Ghavamian, MD *USA*
N Sadoughi, MD *USA*

MA Shalaby, MD *Egypt*
R Sharifi, MD *USA*
S Tabatabaie, MD *USA*
G Vassem, MD *Germany*
Z Zaidi, MD *Pakistan*

AIMS AND SCOPE

As the official journal of the Urology and Nephrology Research Center (UNRC) and the Iranian Urological Association (IUA), *Urology Journal* is a comprehensive digest of useful information on modern urology. Emphasis is on practical information that reflects the latest diagnostic and treatment techniques. Our objectives are to provide an exceptional source of current and clinically relevant research in the discipline of urology, to reflect the scientific work and progress of our colleagues, and to present the articles in a logical, timely, and concise format that meets the diverse needs of today's urologist.

Urology Journal publishes manuscripts on urology and kidney transplantation, all of which undergo extensive peer review by recognized authorities in the field prior to their acceptance for publication. Accordingly, original articles, case reports, and letters to editor are encouraged.

DISCLAIMER

The statements and opinions expressed in *Urology Journal* reflect solely the views of the author(s) and contributor(s) and not of the opinion of IUA or UNRC. The appearance of advertisements in the Journal does not reflect a warranty, endorsement, or approval of any of the products or services described. The UNRC disclaims responsibility for any injury to persons or property resulting from any ideas or products referred to in the articles or advertisements.

Publisher. UNRC

Copyright. UNRC is the copyright owner of the material published in *Urology Journal*. However, in accordance with *Bethesda Statement on Open Access Publishing*, all works published in *Urology Journal* are open access (see the *Instructions for Authors*).

Journal Subscription. Annual: 100 000 Rls (Iran), US\$ 60 (all other countries). Single copy: 30 000 Rls (Iran), US\$ 20 (all other countries).

To subscribe, please contact the editorial office of the journal.

Manuscript Submission. Please prepare your manuscript according to the *Instructions for Authors* of the journal. You can send your manuscript via mail or online submission.

Editorial Office Address. UNRC, No 103, 9th Boustan St, Pasdaran Ave, Tehran 1666679951, Iran. Tel: +98 21 2259 4204, +98 21 2256 7222, Fax: +98 21 2259 4204, Advertising inquiries: +98 21 2259 4204, Web site: <http://www.uj.unrc.ir/>, E-mail: urol_j@unrc.ir

The *Urology Journal* is accredited by the Medical Journals Commission of the Ministry of Health and Medical Education of Islamic Republic of Iran.

Indexed in SCI Expanded, MEDLINE/PubMed, CAB International, Cinahl Database, Cambridge Scientific Abstracts (CSA), Directory of Open Access Journals (DOAJ), Index Copernicus, Google Scholar, and Index Medicus for the WHO Eastern Mediterranean Region (IMEMR)

Front Cover: Top Left, Laparoscopic extraperitoneal simple prostatectomy for benign prostate hyperplasia. See page 107. Top Right, Double ureter and duplex system. See page 145. Bottom, Ureteral herniation with intermittent obstructive uropathy in a renal allograft recipient. See page 98.

REVIEW

Elderly and Prostate Cancer Screening
Stamatiou KN83

SPECIAL FEATURE

Evidence-Based Urology: How Does a Randomized Clinical Trial Achieve Its Designed Goals?
Sadeghi Bazargani H, Hajebrahimi S88

PICTORIAL UROLOGY

Bladder Amyloidosis Mimicking Carcinoma
Singh AK, Floyd MS Jr, De Bolla AR.....97

Ureteral Herniation With Intermittent Obstructive Uropathy in a Renal Allograft Recipient
Mukha RP, Devasia A, Thomas EM.....98

ENDOUROLOGY AND STONE DISEASE

The Most Important Metabolic Risk Factors in Recurrent Urinary Stone Formers
Parvin M, Shakhssalim N, Basiri A, Miladipour AH, Golestan B, Mohammadi Torbati P,
Azadvari M, Eftekhari S99

LAPAROSCOPIC UROLOGY

Laparoscopic Extraperitoneal Simple Prostatectomy for Benign Prostate Hyperplasia: A Two-Year
Experience
Oktay B, Koc G, Vuruskan H, Danisoglu ME, Kordan Y107

UROLOGICAL ONCOLOGY

Bowel Preparation and Peri-operative Management for Radical Cystectomy in Turkey: Turkish
Urooncology Association Multicenter Survey
Aslan G, Baltaci S, Cal C, Turkeri L, Gunlusoy B, Adsan O, and member participants113

Is Positron Emission Tomography Reliable to Predict Post-Chemotherapy Retroperitoneal Lymph
Node Involvement in Advanced Germ Cell Tumors of the Testis?
Akbulut Z, Canda AE, Atmaca AF, Caglayan A, Asil E, Balbay MD120

SEXUAL DYSFUNCTION AND INFERTILITY

General Health and Quality of Life in Patients With Sexual Dysfunctions
Naeinian MR, Shaeiri MR, Hosseini FS127

RECONSTRUCTIVE SURGERY

Triamcinolone Injection Following Internal Urethrotomy for Treatment of Urethral Stricture
Tavakkoli Tabassi K, Yarmohamadi AA, Mohammadi S.....132

General Health and Quality of Life in Patients With Sexual Dysfunctions

Mohammad Reza Naeinian, Mohammad Reza Shaeiri, Fahimeh Sadat Hosseini

Purpose: To study the general health and quality of life in patients with sexual dysfunctions.

Materials and Methods: One hundred and thirty-seven patients with diagnosis of a known sexual dysfunction (SD) were studied. A healthy group of 111 individuals matched for sex, education, and marital status were also selected as a control group. Both groups completed two questionnaires: General Health Questionnaire-28 (GHQ-28) and Personal Wellbeing Index-Adult (PWI-A). To analyze data, descriptive methods as well as student *t* test for independent groups were used.

Results: The mean scores for individuals suffering from SD were more than the control group in total GHQ scale and all its subscales. The mean scores in total PWI-A scale and most of its subscales for individuals suffering from SD were lower than the control group. Since the obtained *t* values (4.16 to 5.22) for all the comparisons done between the mean scores in GHQ for the two groups were higher than *t* value in the 't table' for *df* = 206 at $\alpha = 0.01$ (2.58), differences obtained were significant. Since obtained *t* values (-2.03 to 4.65) for total quality of life and health, achievements, personal relationship, safety, and feeling part of community dimensions were higher than *t* value in the 't table' for *df* = 206 at $\alpha = 0.05$ and $\alpha = 0.01$ (1.96 and 2.58, respectively), differences obtained except for standard of living and future security were significant.

Conclusion: Somatic, social, and mental measures for people having sexual dysfunctions (patient group) were lower than the control group.

Keywords: quality of life, sexual dysfunctions, health status, health surveys

Urol J. 2011;8:127-31.
www.uj.unrc.ir

INTRODUCTION

Sexual problems are common in most of the populations and depending on cultural norms, they surface intermittently in the family practice setting.⁽¹⁾ Sexual dysfunction (SD) is an issue of growing interest. In a population-based study in Iran, of 2626 women interviewed, 31.5% (759) reported SD. The prevalence increased with age from 26% in women aged between 20 and 39 years to 39% in those > 50 years (tested for trend

$P < .001$).⁽²⁾ In another population-based study in Iran, to explore the prevalence of and risk factors for erectile dysfunction (ED), a total of 2674 men aged between 20 and 70 years were interviewed.⁽³⁾ Of the men interviewed, 18.8% (460) reported ED. The prevalence increased with age, from 6% in men aged between 20 to 39 years to 47% in those > 60 years (tested for trend $P < .001$).

Research examining the occurrence

Department of Psychology, Faculty of Arts and Humanity, Shahed University, Tehran, Iran

Corresponding Author:
Mohammad Reza Naeinian, PhD
Department of Psychology, Faculty of Arts and Humanity, Shahed University, Khalij-e-Fars High Way, Tehran, Iran
Tel: + 98 912 770 2588
Fax: + 98 21 5121 2418
E-mail: mnainian@yahoo.com

Received May 2010
Accepted October 2010

of sexual problems in nonclinical populations tends to be restricted to highly selected populations,⁽⁴⁾ such as healthy women in an outpatient gynecological clinic,⁽⁵⁾ normal married couples,⁽⁶⁾ young married couples with children,⁽⁷⁾ and middle-aged men⁽⁸⁾ and women with sexual dysfunction,⁽⁹⁾ with sample size of 38 to 439 subjects.

A review of 23 "community samples" reported a frequency of 4% to 10% for difficulty in achieving orgasm in both men and women, 4% to 9% for erectile problems in men, and 36% to 38% for premature ejaculation in men.⁽⁷⁾ Similarly, a large-scale international collaboration of multidisciplinary experts reported that 40% to 45% of adult women and 20% to 30% of adult men suffer from at least one form of SD. The following prevalence rates were also reported in women: low levels of sexual interest in 17% to 55%, lubrication difficulties in about 8% to 15%, orgasmic dysfunction in 25%, and vaginismus in approximately 6%. The prevalence of ED was reported to be 1% to 9% in men younger than 40 years, which rapidly increased with age to 20% to 40% in men in the age range of 60 to 69 years.⁽¹⁰⁾

It is difficult to obtain an accurate estimate of the prevalence of SD from the international literature because of the discrepancies existing in definitions and tools used in different studies. Only somatic dysfunctions are well-defined, while predominantly psychologically conditioned dysfunctions appear under a multiplicity of labels in various investigations. There is clinical evidence that sexual problems have a multifactorial etiology, including organic, social, and psychological components.⁽¹¹⁾ The impact of certain pathologies, such as depression and diabetes mellitus, on sexual function is well-known.^(12,13) In men, ED is associated with age and is more prevalent in patients suffering from other medical problems.⁽¹⁴⁾ Sexual dysfunctions often coexist with other problems, such as depression, lack of self-esteem, unsuccessful relationships, or just inadequate sexual experience. Nevertheless, very little is known about the relationship between sexual problems and the quality of life.⁽¹⁵⁾

MATERIALS AND METHODS

Target populations were all the people referring to Family Health Clinic in Tehran, with the complaint of a sexual problem. One hundred and thirty-seven patients without a history of other psychiatric disorders were selected for the study by consecutive sampling. They confirmed experiencing a SD through clinical interview by a psychologist, a psychiatrist, or a urologist on the basis of Diagnostic and Statistical Manual, 4th Edition, Text Revision (DSM-IV-TR). One hundred and eleven normal individuals were selected from general population as a control group matched for sex, education, and marital status without having history of sexual problems, to make comparisons possible.

To measure the study outcomes, following instruments were used: 1) Clinical interview on the basis of DSM-IV-TR; 2) General Health Questionnaire-28 (GHQ-28) developed originally by Goldberg⁽¹⁶⁾ and translated into Persian by Taqhavi.⁽¹⁷⁾ Taqhavi reported good psychometric measures (reliability and validity) for the test in Iranian population; 3) Personal Wellbeing Index-Adult (PWI-A), developed by Cummins,⁽¹⁸⁾ is claimed to measure quality of life for adults. Its psychometric properties were confirmed in original articles. Naeinian and colleagues found good psychometric reliability and validity for this tool in Iranian population.⁽¹⁹⁾ Both patient and control groups, who met inclusion criteria for the present study, were individually given the above-mentioned tools initially before starting the treatment.

RESULTS

The patient group consisted of 95 (69.30%) men and 42 (30.70%) women, with the mean age of 49.01 (\pm 12.62) years, while in the control group, 75 (67%) of the participants were men and 36 (32.40%) were women, with the mean age of 40.86 (\pm 12.92) years. Single and married participants in the patient group were 14 (10.20%) and 123 (89.80%), and in the control group were 9 (8.10%) and 102 (91.90%), respectively.

Frequency distribution and percentages of common sexual problems among respondents

are shown in Table 1. Results show that the most common sexual problems were rapid ejaculation in men (27%), reduced sexual desire (21.90%) and vaginismus (15.30%) in women, and performance anxiety (6.6%) and premature erection in men (6.6%).

Descriptive measures, such as mean scores, standard deviations, maximum and minimum scores in GHQ-28 for patients and controls are given in Table 2. Data show that the mean scores for individuals suffering from SD (patient group) were more than the control group in total GHQ scale and all its subscales. As Table 3 shows, the mean scores in total PWI-A scale and most of its subscales for individuals suffering from SD (patient group) were lower than the control group.

Table 1. Frequency distribution and percentage of sexual problems

Diagnosis	Frequency	Percentage (%)
Masturbation	4	2.90
Reduced desire	30	21.90
Vaginismus	21	15.30
Rapid ejaculation	37	27.00
Homosexuality	2	1.50
Performance anxiety	9	6.60
Pain during intercourse	1	0.70
Lack of orgasm	4	2.90
Transvestitism	1	0.70
Premature erection	9	6.60
Lack of pleasure	4	2.90
Frigidity	1	0.70
Sexual aversion	2	1.50
More than one complaint	11	8.10
Unknown	1	0.70
Total	137	100.00

On the basis of data depicted in Table 4, since obtained *t* values (4.16 to 5.22) for all the comparisons done between the mean scores for the two groups were higher than *t* value in the 't table' for *df* = 206 at $\alpha = 0.01$ (2.58), differences obtained were significant. Therefore, general health measures in all studied dimensions were lower in patients suffering from SD in comparison with the control group.

According to Table 5, since obtained *t* values (-2.03 to 4.65) for total quality of life and health, achievements, personal relationship, safety, and feeling part of community dimensions were higher than *t* value in the 't table' for *df* = 206 at $\alpha = 0.05$ and $\alpha = 0.01$ (1.96 and 2.58, respectively), differences obtained except for standard of living and future security were significant. Therefore, total quality of life measure as well as quality of life measure in studied dimensions were lower in patients suffering from SD in comparison with the control group.

DISCUSSION

The most prevalent sexual problems in the studied sample were primary ejaculation, low libido, erection problems, and vaginismus, which were consistent with findings in previous studies.^(7,10) It must be mentioned that apart from cultural and geographical factors in different countries, a proportion of general population in each country suffers from SD, of whom only a limited number seek help.

Results in this study also showed that somatic, social, and psychological measures of people

Table 2. Descriptive data in general health dimensions as measured by GHQ-28*

Dimensions	Group	Statistics			
		Mean	Standard deviation	Minimum	Maximum
Somatic dimension	Patient	7.54	4.15	1	19
	Control	5.36	3.42	0	15
Anxiety and sleepless	Patient	7.57	4.34	0	20
	Control	5.23	3.54	0	18
Social dysfunction	Patient	8.26	2.59	1	19
	Control	6.70	2.55	0	16
Depression	Patient	5.04	5.17	0	21
	Control	2.49	3.24	0	18
Total GHQ score	Patient	28.41	13.65	8	77
	Control	19.78	10.15	1	67

*GHQ-28 indicates General Health Questionnaire-28.

Table 3. Descriptive data in QOL dimensions as measured by PWI-A*

Dimensions	Group	Statistics				
		Mean	Standard deviation	Minimum	Maximum	N
Standard of living	Patient	6	2.13	0	10	137
	Control	4.49	2.42	0	10	111
Health	Patient	5.98	2.46	0	10	137
	Control	7.40	2.29	0	10	111
Achievements	Patient	5.93	2.15	0	10	137
	Control	6.82	2.60	0	10	111
personal relationships	Patient	6.60	2.17	0	10	137
	Control	7.49	1.99	1	10	111
Safety	Patient	6.65	2.57	0	10	137
	Control	7.31	2.48	0	10	111
Feeling part of your community	Patient	6.19	2.38	0	10	137
	Control	6.85	2.36	0	10	111
Future security	Patient	5.85	2.42	0	10	137
	Control	6.02	2.93	0	10	111
Total QOL score	Patient	43.19	12.69	0	69	137
	Control	48.37	13.02	3	70	111

*QOL indicates quality of life; and PWI-A, Personal Wellbeing Index-Adult.

Table 4. Comparison between patients and controls' mean scores in general health dimensions on the basis of student *t* test for independent groups.

Dimensions	Group	Statistics				
		Mean difference	Standard error difference	df	<i>t</i>	<i>P</i>
Somatic dimension	Patient	2.18	0.53	206	4.16	.000
	Control					
Anxiety and sleeplessness	Patient	2.34	0.55	206	4.28	.000
	Control					
Social dysfunction	Patient	1.56	0.36	206	4.38	.000
	Control					
Depression	Patient	2.60	0.59	206	4.32	.000
	Control					
Total GHQ* score	Patient	8.63	1.69	206	5.22	.000
	Control					

*GHQ-28 indicates General Health Questionnaire-28.

Table 5. Comparison between patients and controls' mean scores in QOL dimensions on the basis of student *t* test for independent groups*

Dimensions	Group	Statistics				
		Mean difference	Standard error difference	df	<i>t</i>	<i>P</i>
Standard of living	Patient	-0.49	0.29	246	-1.70	.09
	Control					
Health	Patient	-1.42	0.30	246	-4.65	.000
	Control					
Achievements	Patient	-0.89	0.31	246	-2.96	.003
	Control					
Personal relationships	Patient	-0.89	0.27	246	-3.32	.001
	Control					
Safety	Patient	-0.66	0.32	246	-2.03	.04
	Control					
Feeling part of your community	Patient	-0.66	0.30	246	-2.16	.03
	Control					
Future security	Patient	-0.17	0.34	246	-0.51	.61
	Control					
Total QOL score	Patient	-5.18	1.64	246	-3.16	.002
	Control					

*QOL indicates quality of life.

having SD were lower in comparison with general population. Depressive symptoms have been reported in individuals with SD in earlier studies.⁽¹⁵⁾ Findings in the present study while confirm such previous results, also suggest that adverse effects of sexual problems go more beyond depression. This study also showed that quality of life for people having SD was lower than the control group. This finding is in accordance with the results observed in other countries.^(15,20)

CONCLUSION

We concluded that low general health and quality of life in people with sexual dysfunction cannot be attributed to sexual problems.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Shahar E, Lederer J, Herz MJ. The use of a self-report questionnaire to assess the frequency of sexual dysfunction in family practice clinics. *Fam Pract*. 1991;8:206-12.
- Safarinejad MR. Female sexual dysfunction in a population-based study in Iran: prevalence and associated risk factors. *Int J Impot Res*. 2006;18:382-95.
- Safarinejad MR. Prevalence and risk factors for erectile dysfunction in a population-based study in Iran. *Int J Impot Res*. 2003;15:246-52.
- Fog E, Køster A, Larsen GK, Garde og Inge Lunde K. Female sexuality in various Danish general population age-cohorts. *Nordisk Sexologi*. 1994.
- Rosen RC, Taylor JF, Leiblum SR, Bachmann GA. Prevalence of sexual dysfunction in women: results of a survey study of 329 women in an outpatient gynecological clinic. *J Sex Marital Ther*. 1993;19:171-88.
- Frank E, Anderson C, Rubinstein D. Frequency of sexual dysfunction in "normal" couples. *N Engl J Med*. 1978;299:111-5.
- Nettelblad P, Uddenberg N. Sexual dysfunction and sexual satisfaction in 58 married Swedish men. *J Psychosom Res*. 1979;23:141-7.
- Solstad K, Hertoft P. Frequency of sexual problems and sexual dysfunction in middle-aged Danish men. *Arch Sex Behav*. 1993;22:51-8.
- Osborn M, Hawton K, Gath D. Sexual dysfunction among middle aged women in the community. *Br Med J (Clin Res Ed)*. 1988;296:959-62.
- Lewis RW, Fugl-Meyer KS, Bosch R, et al. Epidemiology/risk factors of sexual dysfunction. *J Sex Med*. 2004;1:35-9.
- Dunn KM, Croft PR, Hackett GI. Association of sexual problems with social, psychological, and physical problems in men and women: a cross sectional population survey. *J Epidemiol Community Health*. 1999;53:144-8.
- Souetre E, Achard F. [Impact of therapeutics on sex. Value of measurements of quality of life]. *Therapie*. 1993;48:461-4.
- Schiel R, Muller UA. Prevalence of sexual disorders in a selection-free diabetic population (JEVIN). *Diabetes Res Clin Pract*. 1999;44:115-21.
- Benet AE, Melman A. The epidemiology of erectile dysfunction. *Urol Clin North Am*. 1995;22:699-709.
- Ventegodt S. Sex and the quality of life in Denmark. *Arch Sex Behav*. 1998;27:295-307.
- Goldberg DP. The detection of psychiatric illness by questionnaire. *Maudsley Monograph No 21*. London: Oxford University Press; 1972.
- Taghavi SM. To study reliability and validity for General Health Questionnaire-28 (GHQ-28). *J Psychol*. 2001;5:381-98.
- Cummins RA, Eckersley R, Lo SK, Okerstrom E. Australian Unity Wellbeing Index Survey 10. Australian Centre for Quality of Life, Deakin University, Melbourne, Report. 2004;10.
- Naeinian MR, Babapour J, Shaeiri MR, Rostami R. The Effect of Neurofeedback Training on the Decrement of Generalized Anxiety Disorder (GAD) Symptoms and Patients, Quality of life. *J Psychol, University of Tabriz*. 2009;5:175-202.
- Lau JT, Kim JH, Tsui HY. Prevalence of male and female sexual problems, perceptions related to sex and association with quality of life in a Chinese population: a population-based study. *Int J Impot Res*. 2005;17:494-505.