

Original Article

A 3-Year Study Of Demographic Characteristics Of Patients With Urinary Tract Infection, Microbial Etiology, And Susceptibility Of Isolated Bacteria To Antibiotics In Shaheed Mostafa Khomeini Hospital

Horieh Sadari¹, Parviz Owlia¹, Mohammad Reza Jalali Nadoushan²,
Farid Zaeri³, Elaheh Zandieh⁴

1. Department of Microbiology, School of Medicine, Shahed University, Tehran, Iran.
2. Department of Pathology, School of Medicine, Shahed University, Tehran, Iran.
3. Department of Health, School of Medicine, Shahed University, Tehran, Iran.
4. School of Medicine, Shahed University, Tehran, Iran.

ABSTRACT

Background and Objectives: This study was designed as a retrospective study on urine samples during three years in Shaheed Mostafa Khomeini Hospital to determine demographic characteristics of patients with urinary tract infection (UTI), microbial etiology, and susceptibility of isolated bacteria to antibiotics.

Materials and Methods: All urines fulfilling the criteria for significant bacteriuria ($>10^4$ colony-forming units/ml of urine) were included in the study. Isolation and identification of bacteria was performed by standard method and susceptibility testing was determined by disk diffusion method according to NCCLS guideline. A total of 909 patients with urinary tract infection were enrolled in this study.

Results: Mean age of the patients was 53.2 years. In addition, females were affected more often than males (female/male sex ratio was 2.22). Meanwhile, considering all strains, 79.5% were Gram-negative bacilli and 67.7% were Enterobacteriaceae. Furthermore, E.coli and Klebsiella spp represented the most common Gram-negative and Enterococci and S. aureus represented the most frequent Gram-positive isolates. The four most frequently isolated bacteria were E. coli (52.1%), Enterococci (10.5%), klebsiella spp. (10.3%), and pseudomonas spp. (9.4%). In addition, E. coli was significantly more common in females (56.6%) than in males (42.2%) and in outpatients (57.4%) than in inpatients (47.4%). The proportion of pseudomonas spp. was significantly higher in males (17.7%) than in females (5.6%). Enterococci were significantly more common in inpatients (12.5%) than in outpatients (8.4%). Altogether, the rate of susceptibility of all UTI pathogens was very low to ampicillin (6.9%) and high to cefotaxime (83.6%) and ciprofloxacin (78.2%). Urinary pathogens isolated from female patients and outpatients were more susceptible to most of examined antibiotics than those isolated from males and inpatients.

Conclusion: It was found out that degrees for antibiotic resistance of urinary pathogens are alarming and show the necessity of keeping up the monitoring of antibiotics susceptibility in UTI isolates and restricting antibiotic consumption in our population.

Keywords: Urinary tract infections, Demographic characteristics, Microbial etiology, Susceptibility

Received: 23 May 2006

Accepted: 20 June 2006

Address communications to: DR. Horieh Sadari, Dept. Microbiology, School of Medicine, Shahed University, Tehran-IRAN.

Email: saderih@yahoo.com.

Introduction

Urinary tract infections (UTI) are amongst the most common infections, with an estimated annual global incidence of at least 250 million (1). It also account for a large proportion of antibiotic consumption (2). Most cases of UTI are initially treated empirically (3). For optimizing the treatment and finding updated recommendations for UTI treatment, physicians should know the etiology of UTI and the susceptibility pattern of UTI pathogens in their population (4). The etiology of UTI and the antimicrobial susceptibility of urinary pathogens have been changing over the years (5). In fact, uropathogens have shown a slow but steady increase in resistance to several antibiotics, for example, *E. coli* and other Enterobacteriaceae have become less susceptible to ampicillin, co-trimoxazole and in some geographical areas, to fluoroquinolones (6). Therefore, this study was performed to determine demographic characteristics of patients with UTI, microbial etiology, and susceptibility of isolated bacteria to commonly used antibiotics during 3 years at Shaheed Mostafa Khomeini Hospital.

Materials and Methods

This study was designed as retrospective and its population consisted of patients with UTI being treated in inpatient section or outpatient clinic of the hospital from 1 January 2001 until 31 December 2003. Demographic data such as age, sex, and type of admission (outpatient or inpatient) were obtained. In addition, UTI was defined as the growth of a single pathogen of $>10^4$ colony forming units/ ml and presence of pyuria in properly collected urine specimen. Identification of isolated bacteria was carried out by standard methods (7).

The antimicrobial susceptibility of bacteria was determined by disk diffusion method using antibiotic-containing disks (Padtan Teb Co., Iran) on Mueller-Hinton agar plate (Merck, Germany) according to guidelines of the NCCLS (8). Susceptibility was tested for ampicillin, carbenicillin, cephalixin, cefazoline, cefixime, ceftazidime, cefotaxime, ceftriaxone, tetracycline, doxycycline, amikacin, gentamicin, kanamycin, ciprofloxacin, ofloxacin, nalidixic acid, nitrofurantoin, and co-trimoxazole

for Gram-negative and to penicillin, ampicillin, oxacillin, vancomycin, cephalixin, cefotaxime, tetracycline, amikacin, gentamicin, kanamycin, ciprofloxacin, nitrofurantoin, co-trimoxazole, erythromycin, and clindamycin for Gram-positive bacteria.

Statistical analysis was performed using SPSS software (version 10) and Pearson's χ^2 as well as Fisher's exact tests, ANOVA, and the Mann-Whitney U tests were applied. A p value less than 0.05 was considered as statistically significant.

Results

A total of 909 non-duplicate bacteria were collected from 627 women (69%) and 282 men (31%) with UTI during the study period. It was found out that females are affected more often than males (female/ male sex ratio was 2.22). In addition, mean age of patients was 53.2 years (53 for females and 53.6 for males). The frequency of pathogens causing UTI is listed in Table 1. Of all studied strains, 79.5% were Gram-negative bacilli and 67.7% were Enterobacteriaceae. The four most frequently isolated bacteria were *E. coli* (52.1%), Enterococci (10.5%), *Klebsiella* spp. (10.3%), and *Pseudomonas* spp. (9.4%). Meanwhile, distribution pattern of isolated bacteria was not significantly different between the years of the study, for example, *E. coli* was identified in 49.1, 51.6, and 54.5% of cases in years 2001, 2002, and 2003 respectively. Furthermore, in patients with UTI due to four most frequently isolated bacteria had the same mean age (52.9, 53.6, 51.1, and 54

Table 1. Distribution of isolated bacteria from UTI

Bacteria	No. of isolates (%)
<i>E. coli</i>	474 (52.1)
Enterococci	96 (10.5)
<i>Klebsiella</i> spp.	94 (10.3)
<i>Pseudomonas</i> spp.	85 (9.4)
<i>S. aureus</i>	48 (5.3)
Enterobacter spp.	36 (4.0)
Acinetobacter spp.	23 (2.5)
Other Streptococci	14 (1.6)
Coagulase negative Staphylococci	14 (1.6)
<i>Streptococcus agalactiae</i>	13 (1.4)
<i>Proteus</i> spp.	12 (1.3)
Total	909 (100)

years for *E. coli*, Enterococci, *Klebsiella* spp. and *Pseudomonas* spp. respectively). Meanwhile, distribution of four most frequently isolated bacteria from male and female cases is shown in Table 2. Predominant pathogen in urine samples of both male and female patients was *E. coli* that significantly occurred more frequently in urines from female (56.6%) than male (42.2%) cases.

Table 2. Distribution of isolated bacteria from male and female cases

Bacteria	No. of isolated bacteria (%)	
	female	male
<i>E. coli</i>	355 (56.6)	119 (42.2)*
Enterococci	59 (9.4)	37 (13.1)
<i>Klebsiella</i> spp.	64 (10.2)	30 (10.6)
<i>Pseudomonas</i> spp.	35 (5.6)	50 (17.7)*
Others	114 (18.2)	46 (16.3)
Total	627 (100)	282 (100)

* A statistically significant difference

In addition, the proportion of *Pseudomonas* spp. was significantly higher in males (17.7%) than in females (5.6%). Bacterial etiology in urines from

inpatients and outpatients with UTI were similar (Table 3). Predominant pathogen in urine samples of both inpatients and outpatients was *E. coli* that significantly occurred more frequently in urines from outpatients (57.4%) than from inpatients (47.4%). The proportion of Enterococci was significantly higher in inpatients (12.5%) than in outpatients (8.4%).

Table 3. Distribution of isolated bacteria from inpatients and outpatients

Bacteria	No. (%) of isolated bacteria from	
	outpatients	inpatients
<i>E. coli</i>	247 (57.4)	227 (47.4) *
Enterococci	36 (8.4)	60 (12.5) *
<i>Klebsiella</i> spp.	47 (10.9)	47 (9.8)
<i>Pseudomonas</i> spp.	32 (7.4)	53 (11.1)
Others	68 (15.8)	92 (19.2)
Total	430 (100)	479 (100)

* A statistically significant difference

On the other hand, the rate of susceptibility to antimicrobial agents is summarized in Tables 4 and 5 for Gram-negative and Gram-positive isolated

Table 4. Susceptibility of Gram-negative isolated bacteria to antibiotics

Antibiotics	Percent of isolates susceptible of :					
	<i>E. coli</i>	<i>Klebsiella</i> spp	Enterobacter spp	Proteus spp	<i>Pseudomonas</i> spp	Acinetobacter spp
Ampicillin	2	0	ND	12.5	0	0
Carbenicillin	33.9	25.6	13.6	ND	18.9	ND
Cephalexin	28.9	46.3	13.3	33.3	ND	ND
Cefazoline	22.1	ND	ND	ND	ND	ND
Cefixime	30.4	39.5	ND	ND	ND	ND
Ceftazidime	71.2	79.3	50	ND	36	ND
Cefotaxime	89	ND	ND	ND	ND	ND
Ceftriaxone	72.7	66.7	ND	ND	3.3	ND
Tetracycline	22.4	33.8	30	ND	18.6	23.5
Doxycycline	12.3	7.7	33.3	ND	10.3	ND
Amikacin	64.7	53.7	39.1	77.8	39.4	14.3
Gentamicin	47	34.4	21.9	50	24.7	14.3
Kanamycin	18	29.6	40	ND	2.3	ND
Ciprofloxacin	79.5	92.3	80	88.9	50	46.7
Ofloxacin	85.7	94.4	ND	ND	34.6	ND
Nalidixic acid	61	57	54.5	58.3	ND	20
Nitrofurantoin	40.8	10.1	27.3	0	1.3	4.8
Co-trimoxazole	29.8	18.9	13.3	25	1.4	9.1

ND = not determined

Table 5. Susceptibility of Gram-positive isolated bacteria to antibiotics

Antibiotics	S. aureus	Percent of isolates susceptible to:			
		Coagulase negative strep.	Enterococci	Strep agalactiae	Other strep
Penicillin	10.3	60	9.9	100	50
Ampicillin	5.9	11.1	27.8	100	58.3
Oxacillin	11.5	ND	ND	ND	ND
Vancomycin	100	100	71.8	100	100
Cephalexin	65.8	92.8	27.7	100	100
Cefotaxime	43.8	ND	43.3	ND	ND
Tetracycline	54.5	54.5	37.3	10	33.3
Amikacin	58.3	90.9	11.3	63.6	60
Gentamicin	51.4	83.3	16.7	30.8	46.2
Kanamycin	23.8	ND	5.6	ND	ND
Ciprofloxacin	74.4	84.6	57.9	90.9	100
Nitrofurantoin	95	76.9	64.6	100	100
Co-trimoxazole	20.5	42.8	25.6	7.7	35.7
Erythromycin	46.3	53.8	34.1	76.9	100
Clindamycin	66.7	50	23.8	90.9	90

ND= not determined

bacteria respectively. In Gram-negative bacilli, the rate of susceptibility to ampicillin was very low and to ofloxacin was higher than others. Vancomycin, nitrofurantoin and ciprofloxacin were more effective against Gram-positive bacteria. In contrast, penicillin and ampicillin had low effect against Gram-positive isolates other than Streptococcus agalactiae. Altogether, the rate of susceptibility of all UTI pathogens was very low to ampicillin (6.9%) and high to cefotaxime (83.6%) and ciprofloxacin (78.2%). Meanwhile, uropathogens isolated from

female patients tends to be come susceptible to most of examined antibiotics than those isolated from males (Table 6) and this difference was statistically significant for some of the antibiotics including ampicillin, cephalexin, cefixime, amikacin, ciprofloxacin, and nitrofurantoin. In addition, antibiotic susceptibility of isolated bacteria from inpatients was lower than outpatients (Table 7) which was statistically significant for cephalexin, cefixime, cefotaxime, gentamicin, ciprofloxacin, nalidixic acid, and nitrofurantoin.

Table 6. Antibiotic susceptibility of bacteria isolated from male and female patients with UTI

Bacteria	Susceptibility rate of isolates from	
	female	male
Ampicillin	8.4	6.2 *
Vancomycin	64.9	60.7
Cephalexin	36	22 *
Cefixime	24.7	22.6 *
Cefotaxime	74.1	65.7
Tetracycline	27	27.1
Amikacin	56.6	46.8 *
Gentamicin	39.8	38
Ciprofloxacin	79.1	67.0 *
Nalidixic acid	54.1	48.8
Nitrofurantoin	47.8	35.9 *
Co-trimoxazole	22.8	26.5

* A statistically significant difference

Table 7. Antibiotic susceptibility rates in bacteria isolated from outpatients and inpatients with UTI

Bacteria	Susceptibility rate of isolates from	
	outpatients	inpatients
Ampicillin	8.7	6.8
Vancomycin	64.1	63.2
Cephalexin	34.7	29.1 *
Cefixime	30.3	19.2 *
Cefotaxime	82	51.8 *
Tetracycline	26	28
Amikacin	54.3	53
Gentamicin	43.5	35.2 *
Ciprofloxacin	77.9	73.1 *
Nalidixic acid	57.6	47.3 *
Nitrofurantoin	46.2	42.3 *
Co-trimoxazole	27.3	20.8

* A statistically significant difference

Discussion

In the present study, an evaluation was conducted on species distribution and antibiotic susceptibility of UTI isolates in Tehran. The most frequent uropathogen was *E. coli* that comprised 52.1% of all isolates in this study. Although a similar frequency for isolates of *E. coli* has been obtained in studies performed in Latin America (52%) (3) and Norway (56.7%) (2), but a different frequency for isolates of *E. coli* has been reported in studies performed in Dakar (38.6%) (9), South Croatia (62.62%) (4), Great Britain (65.1%) (10), and USA (68 and 69% as obtained from two studies) (11, 12). However, in all studied areas, *E. coli* had been the most common pathogen in UTI. According to the demographic data of this study, females were affected more often than males. The question of an underlying urinary pathology of infection in females as compared to male cases was beyond the scope of this study. In addition, the failure rate in female patients with UTI is expected to be greater than in males (13), but isolated bacteria from females in this study tend to be more sensitive to most of the examined antibiotics and this difference was statistically significant for some of them. The same result was obtained in a similar study in Dakar (9). This means that empiric treatment with some of the antibiotics (i.e. ampicillin) is more effective in female than in male cases. The reason for this disparity is not clear. Men may treat themselves empirically, wait a long time before going to clinics, and thus have more resistant bacteria than women in the studied population.

It was also found out that Enterococci caused a significantly higher proportion of infections in inpatients than in outpatients, while *E. coli* was more common in outpatients than in inpatients. This finding correlates with previous studies in Norway (2). In this respect, Enterococci are often multi-resistant and there may be selection for the multi-resistant phenotypes in hospitals, where there is a higher consumption of antibiotics. In fact, in the present study, resistance was higher in isolated bacteria from inpatients than outpatients, which this difference was statistically significant for some of the antibiotics. In addition, the most frequent uropathogen in this study was *E. coli*, which over than 70% of them was resistant to ampicillin, cefalexin, cefazoline, co-trimoxazole, kanamycin,

doxycycline, and tetracycline. High frequency resistance to these antibiotics was also reported among *E. coli* isolates in other studies (2, 3, 9, 14). In contrast, the resistance rate to ceftazidime, cefotaxime, ceftriaxone, ciprofloxacin, and ofloxacin was low in our study which has also been reported in other studies (9, 3). Recommended therapy for UTI is co-trimoxazole, but when rates of resistance to this drug in UTI isolates in population exceeds 10-20%, it is ignored as a first line drug for empirical therapy (4). In the present study, the resistance rate to this antibiotic was 68.4% in 2003, which is much higher than USA and industrial countries, where resistance rate was reported as 29% (4). Therefore, it appears that co-trimoxazole is inadequate in at least two third of cases of urinary tract infections in our population. In this study, the susceptibility rate for UTI isolates to antibiotics was lower than other studies (2-4, 13, 15, 16).

Conclusion

Taken together, it is concluded that degree of antibiotic resistance for uropathogens is alarming and this certainly warrants the necessity of keeping up the monitoring of antibiotics susceptibility in UTI isolates and restricting antibiotic consumption in our population.

Acknowledgements

This study was financially supported by a grant from Shahed University. We also wish to thank laboratory personnel of Shaheed Mostafa Khomeini Hospital for their excellent assistance in data collection.

References

- 1- Ronald AR, Nicolle LE, Stamm E, et al.. Urinary tract infection in adults: research priorities and strategies. *Int J Antimicrob Agents* 2001 17: 343-8.
- 2- Grude N, Tveten Y, Kristiansen BE. Urinary tract infections in Norway: Bacterial etiology and susceptibility. A retrospective study of clinical isolates. *Clin Microbiol Infect* 2001 7: 543-7.
- 3- Gales AC, Jones RN, Gordon KA, et al.. Activity and spectrum of 22 antimicrobial agents tested against urinary tract infection pathogens in hospitalized patients in Latin America: report from the second year of the SENTRY Antimicrobial Surveillance program (1998). *J Antimicrob*

Chemother 2000 45: 295-303.

4- Barisic Z, Babic-Erceg A, Borzic E, et al. Urinary tract infections in South Croatia: aetiology and antimicrobial resistance. *International J Antimicrob Agents* 2003 22: S61-4.

5- Jones RN . Impact of changing pathogens and antimicrobial susceptibility patterns in the treatment of serious infections in hospitalized patients. *Am J Med* 1996 100 (Suppl 6A): S3-12.

6- Garau J, Xercavins M, Rodriguez-Carballeria M, et al. Emergence and dissemination of quinolone-resistant *Escherichia coli* in the community. *Antimicrob Agents Chemother* 1999 43: 2736-41.

7- Baron EJ, Tenover FC, Tenover FC. *Bailey and Scott's Diagnostic Microbiology*, 8th ed. Missouri Mosby Company, 1990.

8- National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial disk susceptibility tests. 8th ed. Approved standard, M2-A8. Wayne, Pennsylvania 2003.

9- Dromigny JA, Nabeth P, Perrier Gros Claude JD. Distribution and susceptibility of bacterial urinary tract infections in Dakar, Senegal. *International J Antimicrob Agents* 2002 20: 339-47.

10- Barrett SP, Savage MA, Rebec MP, et al.. Antibiotic sensitivity of bacteria associated with community-acquired urinary tract infection in Britain. *J Antimicrob Chemother* 1999 44: 359-65.

11- Gupta K, Hooton TM, Wobbe CL, et al. The prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in young women. *Int J Antimicrob Agents* 1999 11: 305-8.

12- Dyer JE, Sankary TM, Dawson JA. Antibiotic resistance in bacterial urinary tract infections, 1991-1997. *West J Med* 1998 169: 265-8.

13- Prais D, Straussberg R, Avitzur Y, et al.. Bacterial susceptibility to oral antibiotics in community acquired urinary tract infection. *Arch Dis child* 2003 88: 215-8.

14- Howard AJ, Magee JT, Fitzgerald KA, Dunstan FDJ. Factors associated with antibiotic resistance in coliform organisms from community urinary tract infection in Wales. *J Antimicrob Chemother* 2001 47: 305- 13.

15- Bouza E, San Juan R, Munoz P, et al.. A European perspective on nosocomial urinary tract infections. I. Report on the microbiology workload, etiology and antimicrobial susceptibility (ESGNI-003 study). *Clin Microbiol Infect* 2001 7: 523-31.

16- Kahlmeter G. The ECO.SENS Project: a prospective, multinational, multicentre epidemiological survey of the

prevalence and antimicrobial susceptibility of urinary tract pathogen- interim report. *J Antimicrob Chemother* 2000 46 (Suppl. S1): 15-22.